























- internal standards including MPA; BMC Clinical Pharmacology,(2012); 12: 1–11.
42. Meinitzer, A., Gartner, G., Pilz, S., Stettin, M.; Ultra fast liquid chromatography-tandem mass spectrometry routine method for simultaneous determination of cyclosporin A, tacrolimus, sirolimus and everolimus in whole blood using deuterated internal standards for cyclosporin A and everolimus; Therapeutic Drug Monitoring,(2010); 32: 61–66.
  43. Vivek Upadhyay, Vikas Trivedi, Gaurang Shah, Manish Yadav and Pranav S. Shrivastav Rapid and Sensitive UPLC–MS–MS Determination of Tacrolimus in Wistar Rats and Human Blood Journal , M. Fathi, J. Veutheyof Chromatographic Science( 2012);1–9.
  44. Chesnut, S.M., Salisbury, J.J.; The role of UHPLC in pharmaceutical development; Journal of Separation Science, (2007); 30:1183–1190.
  45. R.A. Koster, E.C. Dijkers, D.R.Uges, Robust, high-throughput LC-MS/MS method for therapeutic drug monitoring of cyclosporine, tacrolimus, everolimus, and sirolimus in whole blood, Ther. Drug Monit. 31 (2009) 116-125.
  46. M. Karapirli, M. Kizilgun, O. Yesilyurt, H. Gul, Z.I. Kunak, E.O. Akgul, E. Macit, T. Cayci, Y.G. Kurt, I. Aydin, H. Yaren, M. Seyrek, E. Cakir, H. Yaman, Simultaneous determination of cyclosporine A, tacrolimus, sirolimus, and everolimus in whole-blood samples by LC-MS/MS, Scientific World Journal (2012), doi:10.1100/2012/571201
  47. R.A. Koster, J.C. Alffenaar, B. Greijdanus, D.R.A. Uges, Fast LC-MS/MS analysis of tacrolimus, sirolimus, everolimus and cyclosporine in dried blood spots and the influence of the hematocrit and immunosuppressant concentration on recovery, Talanta 115 (2013) 47–54.
  48. B.K. Matuszewski, M.L. Constanzer, and C.M. Chavez-Eng. Strategies for the assessment of matrix effect in quantitative bioanalytical methods based on HPLC-MS/MS. Anal. Chem. 75: 3019-3030 (2003)
  49. M.A. Campaneroa, E. Cardenas, B. Sadaba, E. Garcia-Quetglas, M.J. Munoz-Juarez, I.G. Aldea, D. Pazo, J.R. Azanza, J. Honorato, Therapeutic drug monitoring for sirolimus in whole blood of organ transplants by high-performance liquid chromatography with ultraviolet detection, J. Chromatogr. A, 1031 (2004) 265–273.
  50. M.J. Bogusz, E. Al Enazi, H. Hassan, J. Abdel-Jawaad, J. Al Ruwaily, M. Al Tufail, Simultaneous LC–MS–MS determination of cyclosporine A, tacrolimus, and sirolimus in whole blood as well as mycophenolic acid in plasma using common pretreatment procedure, J. Chromatogr. B 850 (2007) 471–480.
  51. N. Ansermot, J. Desmeules, S. Rudaz, D. Hochstrasser, Simultaneous quantification of cyclosporine, tacrolimus, sirolimus and everolimus in whole blood by liquid chromatography–electrospray mass spectrometry, Clin. Biochem. 41 (2008) 728–735.