











23. Al-Awadi FM, Fatania H, Shamte U. The effect of a plant mixture extract on liver gluconeogenesis in streptozotocin-induced diabetic rats. *Diab. Res.* 1991;18(4):163-168.
24. Mohamed AM, EL-Sharkawy FZ, Ahmed SAA, Aziz WM, Badary OA. Glycemic Control and Therapeutic Effect of *Nigella sativa* and *Curcuma longa* on Rats with streptozotocin-induced Diabetic Hepatopathy. *J. Pharmacol. Toxicol.* 2009;4(2):45-57.
25. Al-Hader A, Aqel M, Hasan Z. Hypoglycemic effect of volatile oil of *Nigella sativa* seeds. *Int. J. Pharmacog.* 1993;31(2):96-100.
26. Farah KM, Atoji Y, Shimizu Y, Takewaki T. Insulinotropic properties of *Nigella sativa* oil in streptozotocin plus nicotinamide diabetic hamsters. *Res. Vet. Sci.* 2002;73:279-282.
27. Matira K, Zesmin FD. Effects of the crude and the n-hexane extract of *Nigella sativa* Linn. (kalajira) upon diabetic rats. *Bangladesh J. Pharmacol.* 2008;4:17-20.
28. Najmi A, Nasiruddin M, Khan RA, Haque SF. Effect of *Nigella sativa* oil on various clinical and biochemical parameters of insulin resistance syndrome. *Int. J. Diab. Dev. Ctries.* 2008;28:11-14.
29. Nadia MH, Taha RA. Effects of *Nigella sativa* Oil and Thymoquinone on Oxidative Stress and Neuropathy in Streptozotocin-Induced Diabetic Rats. *Pharmacology* 2009;84:127-134.
30. Meddah B, Ducroc R, El-Abbes-Faouzi M, Eto B, Mahraoui L, Benhaddou-Andaloussi A, Martineau LC, Cherrah Y and Haddad PS, *Nigella sativa* inhibits intestinal glucose absorption and improves glucose tolerance in rats, *J Ethnopharmacol*, 2009, 21(3), 419-424.
31. Kanter M, Effects of *Nigella sativa* and its major constituent, thymoquinone on sciatic nerves in experimental diabetic neuropathy, *Neurochem Res*, 2008, 33(1), 87-96.
32. Kaleem M, Kirmani D, Asif M, Ahmed Q and Bano B, Biochemical effects of *Nigella sativa* L seeds in diabetic rats, *Indian J Exp Biol*, 2006, 44(9), 745-748.
33. Kanter M, Meral I, Yener Z, Ozbek H and Demir H, Partial regeneration/proliferation of the beta-cells in the islets of Langerhans by *Nigella sativa* L. in streptozotocin-induced diabetic rats, *Tohoku J Exp Med*, 2003, 201(4), 213-219.
34. Kanter M, Coskun O, Korkmaz A and Oter S, Effects of *Nigella sativa* on oxidative stress and beta-cell damage in streptozotocin-induced diabetic rats, *Anat Rec A Discov Mol Cell Evol Biol*, 2004, 279(1), 685-691.
35. Fararh KM, Atoji Y, Shimizu Y, Shiina T, Nikami H and Takewaki T, Mechanisms of the hypoglycaemic and immunopotentiating effects of *Nigella sativa* L. oil in streptozotocin-induced diabetic hamsters, *Res Vet Sci*, 2004, 77(2), 123-129.
36. El-Dakhakhny M, Mady N, Lembert N and Ammon HP, The hypoglycemic effect of *Nigella sativa* oil is mediated by extrapancreatic actions, *Planta Med*, 2002, 68(5), 465-466.
37. Kanter M, Akpolat M and Aktas C, Protective effects of the volatile oil of *Nigella sativa* seeds on beta-cell damage in streptozotocin-induced diabetic rats: a light and electron microscopic study, *J Mol Histol*, 2009, 40(5-6), 379-385.
38. Al-awadi F, Fatima H and Shamte U, The effect of a plant mixture extract on liver gluconeogenesis in streptozotocin induced diabetic rats, *Diabetes Res*, 1991, 18, 163-168.
39. Ebru U, Burak U, Yusuf S, Reyhan B, Arif K, Faruk TH, Emin M, Aydin K, Atilla II, Semsettin S and Kemal E, Cardioprotective effects of *Nigella sativa* oil on cyclosporine A-induced cardiotoxicity in rats, *Basic Clin Pharmacol Toxicol*, 2008, 103(6), 574-580.
40. Burits M and Bucar F, Antioxidant activity of *Nigella sativa* essential oil, *Phytother Res*, 2000, 14(5), 323-328.
41. Badary OA, Thymoquinone attenuates ifosfamide-induced Fanconi syndrome in rats and enhances its antitumor activity in mice, *J Ethnopharmacol*, 1999, 67(2), 135-142.
42. Badary OA, Abdel-Naim AB, Abdel-Wahab MH and Hamada FM, The influence of thymoquinone on doxorubicin-induced hyperlipidemic nephropathy in rats, *Toxicology*, 2000, 143(3), 219-26.
43. Nagi MN, Alam K, Badary OA, Al-Shabanah OA, Al-Sawaf HA and Al-Bekairi AM, Thymoquinone protects against carbon tetrachloride hepatotoxicity in mice via an antioxidant mechanism, *Biochem Mol Biol Int*, 1999, 47(1),153-159.
44. Abuharfeil NM, Salim M and Kleist SV, Augmentation of natural killer cell activity in vivo against tumor cells by wild plants from Jordan, *Phytotherapy Res*, 2001, 15,109-113.
45. Kruk I, Michalska T,Lichszeld K,Kladna A and Aboul- Enein HY, The effect of thymol and its derivatives on reactions generating reactive oxygen species, *Chemosphere*, 2000, 41,1059-1064.
46. Al-Shabanah OA, Badary OA, Nagi MN, Al-Gharably NM, Al-Rikabi AC and Al-Bakairi AM, Thymoquinone protects against doxorubicin induced cardiotoxicity without compromising its antitumor activity, *J Exp Clin Cancer Res*, 1998, 17, 193-198.
47. Nagi M N and Mansour MA, Protective effect of thymoquinone against doxorubicin induced cardiotoxicity in rats: a possible mechanism of protection, *Pharmacol Res*, 2000, 41, 283-289.
48. Badary OA, Al-Shabanah OA, Nagi MN, Al-Rikabi and Elmazar MM, Inhibition of benzopyrene induced forestomach carcinogenesis in mice by thymoquinone, *European J Cancer Prevention*, 1999, 8, 435-440.
49. Badary OA and Gamal-el-Din AM, Inhibitory effects of thymoquinone against 20- methylcholanthrene induce fibrosarcoma tumorigenesis, *Cancer Detec Prev*, 2001, 25, 362-368.
50. Kruk I, Michalska T, Klanda A. The effect of thymol and its derivatives on reaction generating reactive oxygen species. *Chemosphere*. 200;41:1059-1064.
51. Houghton PJ, Zarka R, Heras B, Hoult JR. Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane lipid peroxidation. *Planta Med*. 1995;61:33-36.
52. Badary A, Taha RA, Ayman M, El-Din G, Abdel-Wahab MH. Thymoquinone is a potent superoxide anion scavenger. *Drug Chem. Toxicol.* 2003;26(2):87-98.
53. Daba MH, Abdel-Rehman MS. Hepatoprotective activity of thymoquinone in isolated rat hepatocytes. *Toxicol. Lett.* 1998;95:23-29.
54. Nagi MN, Alam K, Badary OA, Al-Shabanah OA, Al-Sawaf HA, AL- Bekairy AM. Thymoquinone protects against carbon tetrachloride hepatotoxicity in mice via an antioxidant mechanism. *Biochem. Mol. Biol. Int.*1999; 47:153-159.
55. Farah KM, Atoji Y, Shimizu Y, Takewaki T. Insulinotropic properties of *Nigella sativa* oil in streptozotocin plus nicotinamide diabetic hamsters. *Res. Vet. Sci.* 2002;73:279-282.
56. Hamad, Z.M. Protective effect Ethanol extract of *Nigella sativa* L. on hepatic damage induced by naphthalene in male rats. *journal of al-qadisiyah for pure science*, 2012;17 (2):1-10
57. AL-Okaily, B.N.; Mohammed, R.S.; Al-Mzain, K.A. and Khudair, K.K. Effect of Flavonoids Extracted from Black Cumin (*Nigella sativa*) and Vitamin E in Ameliorating Hepatic Damage Induced by Sodium Nitrate in adult male rats. *The Iraqi Journal of Veterinary Medicine*. 2012;36 (2):172-181
58. Mohammed, A.K. Ameliorative effect of black seed (*Nigella sativa* L ) on the toxicity of aluminum in rabbits. *The Iraqi Journal of Veterinary Medicine*.2010;34 (2) :110-116.