



### REFERENCE

- 1) Evaluation and treatment of Cushing's syndrome Lynnette K. Nieman, MD, Ioannis Ilias, MD, DSc Reproductive Biology and Medicine Branch, National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Md. Received: December 31, 2004; Received in revised form: January 28, 2005; Accepted: January 28, 2005;
- 2) LAWRENCE F. KIRK, JR., M.D., Lamar Family Medicine and Occupational Health Center, Barnsville, Georgia ROBERT B. HASH, M.D., HAROLD P. KATNER, M.D., and TOM JONES, M.D., Mercer University School of Medicine, Macon, Georgia Am Fam Physician. 2000 Sep 1;62(5):1119-1127.Cushing's Disease: Clinical Manifestations and Diagnostic Evaluation
- 3) Steffensen C, Bak AM, Rubeck KZ, Jorgensen JO: Epidemiology of Cushing's syndrome. *Neuroendocrinology* 2010, 92(Suppl )
- 4) Jeffcoate WJ, Silverstone JT, Edwards CR, Besser GM: Psychiatric manifestations of Cushing's syndrome: response to lowering of plasma cortisol. *Q J Med* 1979, 48:465-472.
- 5) Nieman LK, Biller BM, Findling JW, Newell-Price J, Savage MO, Stewart PM, Montori VM: The diagnosis of Cushing's syndrome: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Meta*
- 6) Catargi B, Rigalleau V, Poussin A et al. Occult Cushing's syndrome in type 2 diabetes. *J Clin Endocrinol Metab* 2003; 88: 5808-13.
- 7) Howlett TA, Rees LH, Besser GM. Cushing's syndrome. *J Clin Endocrinol Metab* 1985; 14: 911-45.
- 8) Burch WM. *Endocrinology for the house officer*. 3d ed. Baltimore: Williams & Wilkins, 1994:186-9.
- 9) Leibowitz G, Tsur A, Chayen SD et al. Pre-clinical Cushing's syndrome: an unexpected frequent cause of poor glycaemic control in obese diabetic patients. *Clin Endocrinol (Oxf)* 1996; 44: 717-22.
- 10) Pivonello R, Faggiano A, Lombardi G et al. The metabolic syndrome and cardiovascular risk in Cushing's syndrome. *Endocrinol Metab Clin North Am* 2005; 34: 327-39.
- 11) Leibowitz G, Tsur A, Chayen SD, Salameh M, Raz I, Cerasi E, et al. Pre-clinical Cushing's syndrome: an unexpected frequent cause of poor glycaemic control in obese diabetic patients. *Clin Endocrinol (Oxf)*. 1996;44:717-22. [PMID: 8759185]
- 12) Contreras LN, Cardoso E, Lozano MP, Pozzo J, Pagano P, Claus-Hermbeg H. Detección de síndrome de Cushing preclínico en pacientes con sobrepeso y diabetes mellitas tipo 2. *Medicina*. 2000;60:326-30.
- 13) Howlett TA, Rees LH, Besser GM. Cushing's syndrome. *J Clin Endocrinol Metab* 1985; 14: 911-45.
- 14) Biller BM, Grossman AB, Stewart PM, et al. Treatment of adrenocorticotropin-dependent Cushing's syndrome: a consensus statement. *J Clin Endocrinol Metab*. 2008;93(7):2454-2462.