

- 11) I. Alkam T, Nitta A, Mizoguchi H, Saito K, Seshima M, Itoh A *et al.* (2008). Restraining tumor necrosis factor- α by thalidomide prevents the Amyloid β -induced impairment of recognition memory in mice. *Behav Brain Res* 189: 100–106.
- 12) Choi J, Malakowsky CA, Talent JM, Conrad CC, Carroll CA, Weintraub ST *et al.* (2003). Anti-apoptotic proteins are oxidized by Abeta25–35 in Alzheimer’s fibroblasts. *Biochim Biophys Acta* 1637: 135–141.
- 13) R. S. Duman and L. M. Monteggia (2006). “A neurotrophic model for stress-related mood disorders,” *Biological Psychiatry*, vol. 59, no. 12, pp. 1116–1127.
- 14) Sherry L. Xu, Cathy W. C. Bi, Roy C. Y. Choi, Kevin Y. Zhu, Abudureyimu Miernisha, Tina T. X. Dong, and Karl W. K. Tsim (2013). Flavonoids Induce the Synthesis and Secretion of Neurotrophic Factors in Cultured Rat Astrocytes: A Signaling Response Mediated by Estrogen Receptor. Evidence-Based Complementary and Alternative Medicine. Volume 2013, Article ID 127075, 10 pages.
- 15) Tohda C, Tamura T, Komatsu K (2003). Repair of amyloid β (25–35)- induced memory impairment and synaptic loss by a Kampo formula, Zokumei-to. *Brain Res* ,990:141-147.
- 16) Gacar N¹, Mutlu O, Utkan T, Komsuoglu Celikyurt I, Gocmez SS, Ulak G (2011). Beneficial effects of resveratrol on scopolamine butnot mecamlami ne induced memory impairment in the passive avoidance and Morris water maze tests in rats. *Pharmacol Biochem Behav* 99(3):316-23. doi: 10.1016/j.pbb.2011.05.017. Epub May 23.
- 17) Baluchnejadmojarad T, Roghani M, Nadoushan MR, Bagheri M (2009). Neuroprotective effect of genistein in 6-hydroxydopamine Hemi-parkinsonian rat model. *Phytotherapy Research* 23(1):132–135.
- 18) Cain DP (1998). Testing the NMDA, long-term potentiation and cholinergic hypotheses of spatial learning. *Neurosci Biobehav Rev* 22: 181–193.
- 19) M. Parle, D. Dhingra and S. K. Kulkarni (2004). “Neurochemical Basis of Learning and Memory,” *Indian Journal of Pharmaceutical Sciences*, Vol. 66, No. 4, 2004, pp. 371-376.
- 20) A. D. Baddeley (1992). “Working Memory,” *Science*, Vol. 255, No. 5044, 1992, pp. 556-559. <http://dx.doi.org/10.1126/science.1736359>.
- 21) Whitehouse, P. J., Price, D. L., Clark, A. W., Coyle, J. T., & DeLong, M. R. (1981). Alzheimer disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol*, 10(2): 122-126.
- 22) Christine Boesch-Saadatmandia, Agnieszka Lobodab, Anika E. Wagnera, Anna Stachurskab, Alicja Jozkowiczb, Jozef Dulakb, Frank Döringa, Siegfried Wolfframc, Gerald Rimbacha (2011). Effect of quercetin and its metabolites isorhamnetin and quercetin-3-glucuronide on inflammatory gene expression: role of miR-155 ☆ *Journal of Nutritional Biochemistry* 22:293–299.
- 23) J. Kleijnen and P. Knipschild (1992). “Ginkgo biloba for cerebral insufficiency,” *British Journal of Clinical Pharmacology*, vol. 34, no. 4, pp. 352–358, 1992.
- 24) Y. Hou, M. A. Aboukhatwa, D. L. Lei, K. Manaye, I. Khan, and Y. Luo (2010). “Anti-depressant natural flavonols modulate BDNF and beta amyloid in neurons and hippocampus of double TgAD mice,” *Neuropharmacology*, vol. 58, no. 6, pp. 911–920.
- 25) M. Mogi, A. Togari, T. Kondo *et al.* (1999). “Brain-derived growth factor and nerve growth factor concentrations are decreased in the substantia nigra in Parkinson’s disease,” *Neuroscience Letters*, vol. 270, no. 1, pp. 45–48.
- 26) M. Narisawa-Saito, K. Wakabayashi, S. Tsuji, H. Takahashi, and H. Nawa. (1996). “Regional specificity of alterations in NGF, BDNF and NT-3 levels in Alzheimer’s disease,” *NeuroReport*, vol. 7, no. 18, pp. 2925–2928.
- 27) *Psychol Bull.* 1976 May; 83(3):482-504. The Open-Field Test: a critical review. Walsh RN, Cummins RA.