

# A New Cytotoxic and Antioxidant Amentoflavone Monoglucoside from *Cycas revoluta* Thunb Growing in Egypt

Walaa A. Negm, Abd El-Rahim S. Ibrahim, Kamilia A. Abo El-Seoud, Ghada I. Attia, Amany E. Ragab\*  
Department of Pharmacognosy, Faculty of Pharmacy, Tanta university, Tanta 31527, Egypt

## Abstract

Normal and reversed phase silica gel together with Sephadex LH<sub>20</sub> column chromatography of *Cycas revoluta* Thunb extracts of the leaflets led to the isolation and identification of eight compounds (1-8). Compound 6 is a new amentoflavone glucoside to be reported. Compounds 1, 7 and 8 were reported for the first time from *Cycas* genus. Compound 2 was isolated for the first time from *C. revoluta* Thunb. Strikingly, compounds 5, 6 and 7 showed antioxidant activity nearly two to four folds higher than that of quercetin at a dose of 12.5 µg/ml. Compounds 6 and 7 showed very strong cytotoxic activity against MCF 7 cell line (IC<sub>50</sub> = 6.12, 4.73 µg/ml, respectively) compared to doxorubicin (IC<sub>50</sub> = 4.13 µg/ml), while compound 5 showed strong cytotoxic activity (IC<sub>50</sub> = 18.70 µg/ml). The cytotoxic and antioxidant activities are most likely due to the biflavonoid content, especially the new compound 6.

**Key Words:** amentoflavone-4'-O- α- D- glucoside, antioxidant activity, *Cycas revoluta* Thunb, cytotoxic activity.

## INTRODUCTION

Cycadaceae is a family rich in flavonoids, flavonoid glycosides and biflavonoids<sup>[1]</sup>. The only recognized genus of this family is *Cycas* which includes 100 species of which *Cycas revoluta* is the most widely cultivated<sup>[2]</sup>. *Cycas revoluta*, known as Sago palm, is an ornamental plant native to Southern East Asia<sup>[3, 4]</sup> and exhibits cytotoxic, antioxidant, antimicrobial and antileishmanial activities<sup>[5, 6, 7]</sup>. The seeds of *C. revoluta* are used in Chinese medicine as expectorant and antirheumatic<sup>[8]</sup>. Phytochemical investigation of the plant leaflets in previous studies led to the isolation of flavonoids (naringenin;<sup>[6]</sup> prunin;<sup>[5]</sup> vitexin-2''-rhamnoside<sup>[5]</sup>), biflavonoids (amentoflavone;<sup>[9]</sup> 2,3-dihydroamentoflavone;<sup>[6]</sup> 2, 3, 2'', 3''-tetrahydroamentoflavone; 2, 3-dihydro- 7- O- β- D-glucopyranosylamentoflavone;<sup>[5]</sup> 2, 3- dihydro- 7, 7''- di-O- β-D-glucopyranosylamentoflavone;<sup>[5]</sup> hinokiflavone;<sup>[6]</sup> 2, 3-dihydrohinokiflavone;<sup>[9]</sup> 2, 3, 2'', 3''-tetrahydrohinokiflavone;<sup>[6]</sup> podocarpusflavone A),<sup>[6]</sup> phenolic and alcoholic compounds ((+)-lariciresinol;<sup>[6]</sup> protocatechuic acid;<sup>[5]</sup> vomifoliol<sup>[6]</sup>). Little data were found in literature regarding the cytotoxic activity and no data were found about the antioxidant activity of the active principles from *C. revoluta* leaflets<sup>[5, 6]</sup>. Here we report the isolation of one new and four known compounds from *C. revoluta* leaflets for the first time from the genus *Cycas* and explore the antioxidant and cytotoxic activities of the studied extracts and some purified compounds.

## MATERIALS AND METHODS

### GENERAL EXPERIMENTAL PROCEDURES

Authentic β-sitosterol was purchased from Merck, Darmstadt, Germany. Authentic amentoflavone was kindly provided by Prof. Abdel-Rahim Sayed Ibrahim, Professor of Pharmacognosy, Faculty of Pharmacy, Tanta University. Naringenin, anhydrous Na<sub>2</sub>SO<sub>4</sub>, glucose, rhamnose, AlCl<sub>3</sub>, NaOA, and boric acid were obtained from were obtained

from Sigma Chemical Co., USA. Melting points were measured using melting point apparatus "Galenkamp type" and were uncorrected. Thin layer chromatography (TLC) was performed using silica gel G F<sub>254</sub> plates (E. Merk). Column chromatography was accomplished using silica gel (E. Merck, 70-230 mesh). Sephadex LH-20 was purchased from Sigma- Aldrich chemical Co., USA. Reversed phase octadecylsilyl-silica gel (RP-C18) was obtained from Merck, Germany and Diaion™ HP20SS from Mitsubishi Chemical, Japan. Sheets of Whatman No. 1 filter paper for chromatography were obtained from Whatman Ltd., England. NMR spectra were measured using Bruker High Performance Digital FT-NMR Spectrometer Avance III operating at 400 MHz. IR spectra were recorded as KBr discs using Jasco, FT/IR-6100 spectrophotometer, Japan. UV spectra were obtained using Shimadzu UV spectrophotometer UV- 1800, Japan. EI-MS spectra were recorded using mass Spectrometer Thermo Scientific ISQ Single Quadrupole MS, USA and mass Spectrometer Thermo Scientific ISQ Quantum Access MAX triple Quadrupole, USA.

### BIOLOGICAL ACTIVITY

2, 2- Diphenyl-1- picrylhydrazyl (DPPH); dimethyl sulfoxide (DMSO); trichloroacetic acid; sulforhodamine B (SRB) stain; ethylenediaminetetraacetic acid buffer (EDTA) and acetic acid were purchased from Sigma Chemical Co. St. Louis, Mo. Quercetin was purchased from Fluka Chemical Crop, USA. Breast carcinoma cell line (MCF-7) and liver carcinoma cell line (HepG2) were obtained from the American Type Culture Collection (ATCC). Absorbance in the antioxidant assay was measured using a UNICO spectrophotometer (UV/Vis) UV-2000 (UNICO Instrument CO. LTD, USA) and an ELISA processor II microplate reader was used in the cytotoxic activity assessment.

**Plant material**

*Cycas revoluta* Thunb leaflets were collected in August 2011 from local garden in Tanta city, Gharbia Governorate, Egypt. The plant was kindly identified by Prof. Kamal Shaltout, professor of plant ecology, botany department, Faculty of Science, Tanta University. They were shade-dried, pulverized and stored in amber-colored glass bottles for further investigation.

**Extraction and isolation**

The shade dried powder of *Cycas revoluta* Thunb leaves (2 kg) was extracted by cold maceration with ethanol (95%) until exhaustion. Alcohol extract was evaporated under reduced pressure to yield 70 gm of crude total extract, which was resuspended in aqueous methanol (50%) and successively fractionated with petroleum ether (40-60 °C), methylene chloride, ethyl acetate and n- butanol saturated with water.

**Pet. ether extract** (2 g) was saponified by boiling under reflux with 10% alcoholic KOH (30 ml) for 5 hrs. The major part of alcohol was then distilled off and the aqueous liquid left was diluted with 20 ml water and extracted with several portions of ether till exhaustion.<sup>[10]</sup> The combined ethereal extract was washed with distilled water, dehydrated over anhydrous Na<sub>2</sub>SO<sub>4</sub> and distilled off under reduced pressure to give the unsaponifiable matter (0.37 g). The unsaponifiable matter was subjected to column chromatography on silica gel (1.5 x 35 cm, 30 g) gradiently eluted using pet. ether-methylene chloride mixtures of increasing polarity. Fractions (52-61) eluted with pet. ether-methylene chloride (50:50) yielded white powder (90 mg) which was recrystallised from methanol to afford compound **2** as white needle crystals (25 mg). Fractions (20-32) collected with pet. ether- methylene chloride (80:20) were pooled, evaporated to dryness (20 mg) and rechromatographed on silica gel column eluted with methylene chloride- methanol (2% increment) to yield compound **1** (4 mg).

**Methylene chloride fraction** (4 g) of the crude total extract was chromatographed on silica gel (3 x 60 cm, 105 g) gradiently eluted with methylene chloride- methanol mixtures of increased polarity. Fractions (4-10) eluted with methylene chloride- methanol (90:10) were pooled and evaporated to dryness to yield 800 mg and rechromatographed on silica gel column (2.5 x 47 cm, 25 g). The column was eluted with n-hexane- ethyl acetate mixtures of which the 50:50 eluate was further purified on sephadex gel column (1.25 x 25 cm, 10g) to yield compound **3**. Fractions (11-20) eluted with methylene chloride- methanol (86:14) were pooled, evaporated to dryness to yield 360 mg and rechromatographed on silica gel column (1.25 x 25 cm, 15 g) eluted with methylene chloride and increasing polarity using methanol of which the methylene chloride- methanol 94:6 and 90:10 eluates were pooled separately. The methylene chloride- methanol 94:6 eluate (54 mg) was further chromatographed on silica gel mini column (0.7x 10 cm, 2 g) using methylene chloride- methanol (95:5) to afford compound **4** (21 mg). The methylene chloride- methanol 90:10 eluate (110 mg) was further purified on a Sephadex LH<sub>20</sub> gel column eluted with methanol to yield compound **5** (60 mg).

**Ethyl acetate fraction** (2 g) of the crude total extract was chromatographed on a silica gel column (3 x 60 cm, 75 g) gradiently eluted with methylene chloride- methanol mixtures of increasing polarity. Fractions eluted with methylene chloride- methanol (88:12) and (84:16) were pooled separately for further purification. The methylene chloride- methanol 88:12 eluate (fractions 23-30, 174 mg) was rechromatographed on a reversed phase C-18 silica gel column (1.5 x 32 cm, 8 g) eluted with 10% MeOH in water for HPLC and decreasing polarity by HPLC methanol of which the 40:60 eluate provided compound **6** (44 mg). The Fractions 31-40 obtained with methylene chloride- methanol (84:16) of ethyl acetate fraction was rechromatographed on silica gel column (0.7 x 25 cm, 2.5 g) gradiently eluted with methylene chloride- methanol mixtures of increasing polarity of which the 88:12 to 84:16 eluates yielded compound **7** (20 mg) after evaporation to dryness.

**n- Butanol fraction** (7.5 gm) of crude total extract was suspended in deionized water and applied on top of a Diaion HP-20 column (3 x 60 cm, 80 g). The column was initially eluted with deionized water (1 L) to remove free sugars and salts followed by water and methanol mixtures of decreasing polarity of which 60% methanol eluate (500 mg) was rechromatographed on a silica gel column (1.5 x 60 cm, 15 g) and eluted with methylene chloride and increasing polarity using methanol of which the methylene chloride-methanol (75:25) eluate provided compound **8** (16 mg) after repeated crystallization from methanol.

**Acid hydrolysis of the isolated glycosides<sup>[11]</sup>**

Complete acid hydrolysis was carried out by refluxing the glycoside (5 mg) in aqueous methanol with 5% H<sub>2</sub>SO<sub>4</sub> (10 ml) for 2 hrs. The hydrolysate was extracted with ethyl acetate and washed with water, then evaporated to obtain the aglycone. The mother liquor of hydrolysate was neutralized with Ba<sub>2</sub>CO<sub>3</sub>, filtered, concentrated and used to study the sugar moiety using paper chromatography.

**BIOLOGICAL SCREENING****Antioxidant activity**

The antioxidant activity was assessed by measuring the free radical scavenging activity of the different extracts and some isolated pure compounds using DPPH according to literature procedure<sup>[12, 13, 14]</sup>. Tested solutions were prepared in ethanol. The test was carried out, in triplicate, by mixing 0.2 ml of 1 mM DPPH solution, 0.2 ml of tested sample and 4.6 ml ethanol. The mixture was incubated at 30 °C for 30 min in the dark. The absorbance was measured at 517 nm using ethanol as a blank. Quercetin was used as a positive control and negative control was prepared by omitting the addition of test sample. The percent of inhibition of free radical DPPH (I %) was calculated.

**Cytotoxic activity**

Potential Cytotoxicity of the different extracts and isolated pure compounds was investigated according to Skehan *et al*<sup>[15]</sup> using MCF-7 and HepG2 cell lines. Doxorubicin was used as a standard and the IC<sub>50</sub> was calculated and assessed according to Ayyad classification<sup>[16]</sup>.

**CHARACTERIZATION****Compound 1**

White crystal; mp 188-189°C; IR (KBr)  $\nu_{\max}$  3443, 2934, 2865, 1715, 1642, 1460, 1372  $\text{cm}^{-1}$ ; EI-MS  $m/z$  426 [ $\text{M}^+$ ], 411 [ $\text{M}^+ - \text{CH}_3$ ], 408 [ $\text{M}^+ - \text{H}_2\text{O}$ ], 381 [ $\text{M}^+ - 3\text{CH}_3$ ].

**Compound 2**

White crystals; mp 137-139°C; IR (KBr)  $\nu_{\max}$  3449, 2936, 2861, 2359, 2069, 1640, 1461, 1377, 1329, 1244, 1187, 1107, 1056, 961, 924, 877, 833, 800, 619  $\text{cm}^{-1}$ ; EI-MS  $m/z$  414 [ $\text{M}^+$ ], 396 [ $\text{M}^+ - \text{H}_2\text{O}$ ], 381 [ $\text{M}^+ - (\text{CH}_3 + \text{H}_2\text{O})$ ], 329 [ $\text{M}^+ - \text{C}_6\text{H}_{13}$ ];  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 5.26 (1H, t, H-6), 3.50 (1H, m, H-3), 1.24 (3H, s, H-19), 0.90 (3H, d,  $J = 6$  Hz, H-21), 0.84 (3H, d,  $J = 6.4$  Hz, H-26), 0.81 (3H, d,  $J = 6.8$  Hz, H-27), 0.65 (3H, s, H-18), 0.65-2.24 (m, H-1, 2, 4, 7-9, 11, 12, 14-17, 20, 22-25, 28, 29).

**Compound 3**

White amorphous granules; UV  $\lambda_{\max}$  (MeOH) 293, 325 (sh) nm; IR (KBr)  $\nu_{\max}$  3957, 3780, 3290, 3119, 2918, 2831, 2709, 1634  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 5.45 (1H, dd,  $J = 12.8, 3.4$  Hz, H-2), 2.69 (1H, dd,  $J = 17.2, 3.4$  Hz, H-3 axial), 3.30 (1H, dd,  $J = 17.0, 12.8$  Hz, H-3 equatorial), 5.88 (2H, d,  $J = 2.1$  Hz, H-6, 8), 7.31 (2H, d,  $J = 8.0$  Hz, H-2', 6'), 6.81 (2H, d,  $J = 8.0$  Hz, H-3', 5'), 12.15 (1H, s, OH).

**Compound 4**

Yellow amorphous granules; UV  $\lambda_{\max}$  (MeOH) 269, 280, 305 (sh) and 332 nm; IR (KBr)  $\nu_{\max}$  3858, 3450, 2927, 2859, 2353, 2319, 2072, 1644, 1995, 1430, 1364, 1287, 1247, 1170, 1109, 1042, 947, 835, 643, 593  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 6.76 (1H, s, H-3"), 6.81 (1H, s, H-3), 6.18 (1H, d,  $J = 1.8$  Hz, H-6), 6.28 (1H, s, H-6"), 6.41 (1H, d,  $J = 1.8$  Hz, H-8), 7.60 (2H, d,  $J = 8.6$  Hz, H-2"', 6'''), 6.68 (2H, d,  $J = 8.6$  Hz, H-3"', 5'''), 7.98 (1H, dd,  $J = 8.6, 2.3$  Hz, H-6'), 8.08 (1H, d,  $J = 2.3$  Hz, H-2'), 7.07 (1H, d,  $J = 8.6$  Hz, H-5'), 10.77 (1H, s, OH at C7), 10.21 (1H, s, OH at C7"), 13.02 (1H, s, OH at C5), 13.14 (1H, s, OH at C5"); ESI-MS  $m/z$  553 [ $\text{M} + \text{H}$ ] $^+$ .

**Compound 5**

Yellow amorphous powder; UV  $\lambda_{\max}$  (MeOH) 270, 332 nm; IR (KBr)  $\nu_{\max}$  3963, 3426, 1654, 1610, 1500, 1429, 1360, 1285, 1244, 1172, 1108, 1040, 836, 754, 560  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 6.79 (1H, s, H-3"), 6.83 (1H, s, H-3), 6.19 (1H, d,  $J = 1.8$  Hz, H-6), 6.40 (1H, s, H-6"), 6.47 (1H, d,  $J = 1.8$  Hz, H-8), 7.58 (2H, d,  $J = 8.0$  Hz, H-2"', 6'''), 6.73 (2H, d,  $J = 8.0$  Hz, H-3"', 5'''), 7.15 (1H, d,  $J = 8.8$  Hz, H-5'), 8.00 (1H, d,  $J = 1.5$  Hz, H-2'), 7.99 (1H, dd,  $J = 8.8, 1.5$  Hz, H-6'), 10.86 (1H, s, OH at C7), 10.31 (1H, s, OH at C7"), 10.65 (2H, brs, OH at 4' and 4''), 12.97 (1H, s, OH at C5), 13.10 (1H, s, OH at C5");  $^{13}\text{C}$  NMR [DMSO- $d_6$ , 100 MHz]  $\delta_{\text{C}}$ : 164.3 (C-2), 103.4 (C-3), 182.2 (C-4), 161.9 (C-5), 99.1 (C-6), 164.2 (C-7), 94.5 (C-8), 157.8 (C-9), 104.2 (C-10), 120.5 (C-1'), 128.2 (C-2'), 121.9 (C-3'), 160.9 (C-4'), 116.7 (5'), 131.8 (C-6'), 164.6 (C-2''), 103.1 (C-3''), 182.6 (C-4''), 160.1 (C-5''), 99.3 (C-6''), 162.4 (C-7''), 104.5 (C-8''), 154.9 (C-9''), 104.1 (C-10''), 121.4 (C-1'''), 128.6 (C-2''', 6'''), 116.2 (C-3''', 5'''), 161.5 (C-4'''); ESI-MS  $m/z$  539 [ $\text{M} + \text{H}$ ] $^+$ .

**Compound 6**

Yellow amorphous powder; UV  $\lambda_{\max}$  (MeOH) 271, 332 nm; IR (KBr)  $\nu_{\max}$  3928, 3782, 3420, 2926, 2860, 2305, 1723, 1650, 1609, 1498, 1429, 1362, 1285, 1245, 1172, 1106, 948, 836, 744, 637, 560, 513  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 6.80 (1H, s, H-3"), 6.82 (1H, s, H-3), 6.34 (1H, s, H-6"), 6.19 (1H, d,  $J = 2.1$  Hz, H-6), 6.44 (1H, d,  $J = 2.1$  Hz, H-8), 7.6 (2H, d,  $J = 8.4$  Hz, H-2"', 6'''), 6.77 (2H, d,  $J = 8.4$  Hz, H-3"', 5'''), 8.00 (1H, d,  $J = 2.1$  Hz, H-2'), 7.12 (1H, d,  $J = 8.4$  Hz, H-5'), 7.99 (1H, dd,  $J = 8.4, 2.1$  Hz, H-6'), 10.85 (1H, s, OH at C7), 10.30 (1H, s, OH at C7"), 13.11 (1H, s, OH at C5), 12.99 (1H, s, OH at C5"), 4.47 (1H, d,  $J = 4.4$  Hz, anomeric proton);  $^{13}\text{C}$  NMR [DMSO- $d_6$ , 100 MHz]  $\delta_{\text{C}}$ : 164.4 (C-2), 103.0 (C-3), 182.2 (C-4), 161.4 (C-5, 5''), 99.3 (C-6), 161.9 (C-7), 94.5 (C-8), 157.3 (C-9), 104.1 (C-10), 127.8 (C-1'), 128.0 (C-2', 6'), 121.9 (C-3'), 161.0 (C-4'), 121.8 (5'), 164.6 (C-2''), 103.2 (C-3''), 182.5 (C-4''), 99.5 (C-6''), 164.0 (C-7''), 104.1 (C-8''), 154.9 (C-9''), 104.0 (C-10''), 121.9 (C-1'''), 128.6 (C-2''', 6'''), 116.2 (C-3''', 5'''), 161.7 (C-4'''), 99.9 (Glu1), 70.0 (Glu2), 72.9 (Glu3), 67.9 (Glu4), 70.2 (Glu5), 63.5 (Glu6); ESI-MS  $m/z$  701 [ $\text{M} + \text{H}$ ] $^+$ .

**Compound 7**

Yellow amorphous powder; UV  $\lambda_{\max}$  (MeOH) 227, 283, 340 (sh) nm; IR (KBr)  $\nu_{\max}$  3449, 2958, 2042, 1639, 1586, 1506, 1270, 1178, 1128, 1127, 1022, 875, 818, 745, 647, 547  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 5.52 (1H, dd,  $J = 12.6, 3.2$  Hz, H-2), 2.55 (1H, dd,  $J = 16.3, 3.2$  Hz, H-3 axial), 3.78 (1H, dd,  $J = 16.3, 12.6$  Hz, H-3 equatorial), 6.12 (1H, d,  $J = 3.6$  Hz, H-6), 6.14 (1H, d,  $J = 3.6$  Hz, H-8), 6.96 (1H, d,  $J = 1.6$  Hz, H-2'), 6.91 (1H, d,  $J = 8.4$  Hz, H-5'), 6.94 (1H, dd,  $J = 1.6, 8.4$  Hz, H-6'), 3.9 (3H, s,  $\text{OCH}_3$ ), 4.97 (2H, d,  $J = 7.4$  Hz, glucose anomeric proton), 4.69 (H, d,  $J = 2.8$  Hz, rhamnose anomeric proton), 1.09 (3H, d,  $J = 6.2$  Hz,  $\text{CH}_3$  of rhamnose), 9.12 (1H, s, OH at C-3'), 12.01 (1H, s, OH at C-5);  $^{13}\text{C}$  NMR [DMSO- $d_6$ , 100 MHz]  $\delta_{\text{C}}$ : 78.5 (C-2), 40.9 (C-3), 197.5 (C-4), 163.3 (C-5), 96.8 (C-6), 165.6 (C-7), 95.8 (C-8), 162.5 (C-9), 104.6 (C-10), 131.3 (C-1'), 112.5 (C-2'), 146.9 (C-3'), 148.4 (C-4'), 114.6 (C-5'), 118.4 (C-6'), 99.9 (Glu1), 76.7 (Glu2), 76.0 (Glu3), 71.2 (Glu4), 73.5 (Glu5), 59.8 (Glu6), 101.1 (Rha1), 70.1 (Rha 2), 70.7 (Rha 3), 72.5 (Rha 4), 68.8 (Rha5), 18.3 ( $\text{CH}_3$  of Rha 6), 56.2 ( $\text{OCH}_3$ ); ESI-MS  $m/z$  611 [ $\text{M} + \text{H}$ ] $^+$ .

**Compound 8**

Yellow granules; UV  $\lambda_{\max}$  (MeOH) 229, 260, 357 nm; UV  $\lambda_{\max}$  (NaOH) 258, 288, 408; ( $\text{AlCl}_3$ ) 272, 305 (sh), 430; UV  $\lambda_{\max}$  ( $\text{AlCl}_3/\text{HCl}$ ) 302, 370, 400; UV  $\lambda_{\max}$  (NaOAc) 274, 412; UV  $\lambda_{\max}$  (NaOAc/Boric acid) 264, 374; IR (KBr)  $\nu_{\max}$  3419, 2925, 2855, 1627, 1383  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR [DMSO- $d_6$ , 400 MHz]  $\delta_{\text{H}}$ : 6.14 (1H, d,  $J = 2.0$  Hz, H-6), 6.33 (1H, d,  $J = 2.0$  Hz, H-8), 6.83 (1H, d,  $J = 8.4$  Hz, H-5'), 7.54 (2H, m, H-2', 6'), 4.39 (1H, d,  $J = 2.0$  Hz, rhamnose anomeric proton), 5.32 (1H, d,  $J = 7.0$  Hz, glucose anomeric proton), 0.99 (3H, d,  $J = 6.0$  Hz,  $\text{CH}_3$  of rhamnose), 12.58 (1H, s, OH at C-5).

## RESULTS AND DISCUSSION

### PHYTOCHEMICAL INVESTIGATION

$\beta$ - Amyrin (**1**) and  $\beta$ - sitosterol (**2**) were identified by comparing their physical and spectral data to that of authentic samples and published literature<sup>[17-20]</sup>.

#### Compound 3

UV spectrum showed  $\lambda_{\max}$  characteristic for flavanones at 293 and 340 (sh) nm<sup>[21]</sup>. <sup>1</sup>H-NMR data confirmed the flavanone nature of compound **3** as it exhibited signals for methylene protons at C-3 resonating at  $\delta_{\text{H}}$  2.69 (dd,  $J=17.2$ , 3.4 Hz) and at  $\delta_{\text{H}}$  3.3 (dd,  $J=17.0$ , 12.8). Additionally, <sup>1</sup>H NMR spectrum indicated an AA'BB' system by signals at  $\delta_{\text{H}}$  7.31 (d,  $J=8.0$  Hz) for H-2', 6' and 6.81 (d,  $J=8.0$  Hz) for H-3', 5'. A signal at  $\delta_{\text{H}}$  5.88 (2H, d,  $J=2.1$  Hz) is attributed to *meta*-coupled equivalent protons H-6, 8. The UV and <sup>1</sup>H NMR data were found matching those of naringenin in literature<sup>[22,23]</sup>. The structure of compound **3** was confirmed by direct comparison with naringenin authentic sample through TLC and superimposable IR & UV spectra.

#### Compound 4

UV spectrum showed  $\lambda_{\max}$  at 269 and 332 nm, which suggested a flavonoid structure. <sup>1</sup>H NMR data confirmed the biflavonoid nature of compound **4** and suggested an amentoflavone pattern<sup>[24]</sup>. The <sup>1</sup>H NMR spectrum showed an AA'BB' system as indicated by the signal at  $\delta_{\text{H}}$  7.60 (d, 8.6 Hz) for H-2'', 6'' and at  $\delta_{\text{H}}$  6.68 (d, 8.6 Hz) for H-3'', 5''. An ABX system was also shown by the signals at  $\delta_{\text{H}}$  8.08 (d,  $J=2.3$  Hz) for H-2',  $\delta_{\text{H}}$  7.07 (d,  $J=8.6$  Hz) for H-5' and  $\delta_{\text{H}}$  7.98 (dd,  $J=8.6$ , 2.3 Hz) for H-6'. A singlet at  $\delta_{\text{H}}$  6.28 and *meta*-coupled protons at  $\delta_{\text{H}}$  6.18 and 6.41 ( $J=1.8$  Hz) were assigned for H-6'', H6 and H-8, respectively. The NMR data indicated that C-8'' and C-3' are involved in the biflavonoid linkage. The ESI<sup>+</sup>-MS of compound **4** showed a pseudomolecular ion at  $m/z$  553.0 for [M+H]<sup>+</sup> which is consistent with an amentoflavone methoxy derivative. Inspection of the <sup>1</sup>H NMR spectrum revealed the absence of 4', 4'''-OH signal and showed a new signal suggesting 4'-methoxy amentoflavone structure at either site. The UV spectrum of compound **4** in methanol exhibited  $\lambda_{\max}$  at 269, 280, 305 (sh) and 332 nm which is consistent with 4'''- and not 4'- methoxy derivative<sup>[25]</sup>. In the <sup>1</sup>H NMR the methoxy signal was buried under the residual water peak. Based on all the data discussed above, compound **4** was determined as podocarpus flavone A which was previously isolated from the plant<sup>[6]</sup>.

#### Compound 5

UV spectrum showed  $\lambda_{\max}$  at 270 and 332 nm which suggested a flavonoid structure. <sup>1</sup>H-NMR data confirmed the biflavonoid nature of compound **5** and suggested an amentoflavone pattern. The <sup>1</sup>H NMR spectrum showed an AA'BB' systems as indicated by  $\delta_{\text{H}}$  6.72 (d, 8.0 Hz) for H-3''', 5''' and  $\delta_{\text{H}}$  7.58 (d,  $J=8.0$  Hz) for H-2''', 6'''. Additionally, the <sup>1</sup>H NMR spectrum showed signals for *meta*-coupled protons at  $\delta_{\text{H}}$  6.19 and 6.46 ( $J=1.8$  Hz) for H-6 and H-8, respectively, also a singlet at  $\delta_{\text{H}}$  6.40 was assigned to H-6''. An ABX system was also shown by the signals at  $\delta_{\text{H}}$  8.00 (d,  $J=1.5$  Hz) for H-2',  $\delta_{\text{H}}$  7.15 (d,  $J=8.8$  Hz) for H-5' and  $\delta_{\text{H}}$  7.99 (dd,  $J=8.8$ , 1.5 Hz) for H-6'. These data indicated

that C-8'' and C-3' are involved in the biflavonoid linkage which is consistent with amentoflavone in literature. The structure was confirmed by <sup>13</sup>C NMR which showed 30 carbons and confirmed the conjugation linkage between C-8'' and C-3' due to the downfield shift of C-8'' by 10 ppm and of C-3'' by 6.00 ppm, respectively<sup>[26, 27, 28]</sup>. The ESI<sup>+</sup>-MS of compound **5** showed a pseudomolecular ion at  $m/z$  539.34 for [M+H]<sup>+</sup> which is consistent with an amentoflavone structure. The NMR data were consistent with those in literature for amentoflavone<sup>[24, 26-29]</sup>.

#### Compound 6

The results of chemical tests, Molisch's and NaOH tests, of compound **6** suggested a flavonoid glycoside nature. IR spectrum showed a strong band at 3420 cm<sup>-1</sup> for hydroxyl stretching, 1650 cm<sup>-1</sup> for carbonyl group, 2926 cm<sup>-1</sup> for aliphatic methylene stretching, 1362 cm<sup>-1</sup> for methylene bending and 1285 cm<sup>-1</sup> for (C-O) stretching. UV spectrum showed  $\lambda_{\max}$  at 271, 299 and 332 nm which are characteristic of flavonoids<sup>[27]</sup>. <sup>1</sup>H NMR data confirmed the biflavonoid nature of compound **6** and suggested an amentoflavone pattern as compared to compound **5** data discussed above. <sup>1</sup>H NMR spectrum established the presence of two hydroxyl groups resonating at  $\delta_{\text{H}}$  13.11, 12.99 indicating the presence of two chelated hydroxyl groups at C-5, 5'', respectively. The signals of *meta*-coupled protons at  $\delta_{\text{H}}$  6.19 and 6.44 (d,  $J=2.1$  Hz) can be assigned for H-6 and H-8, respectively while the signal at  $\delta_{\text{H}}$  6.34 (1H, s) accounts for H-6''. The signals at  $\delta_{\text{H}}$  6.82 ppm (s) and 6.80 (s) are for H-3 and H-3'', respectively. The <sup>1</sup>H NMR spectrum also showed an ABX system represented by signals at  $\delta_{\text{H}}$  8.00 (d,  $J=2.1$  Hz) for H-2', 7.12 (d,  $J=8.4$  Hz) for H-5', 7.99 (dd,  $J=8.4$ , 2.1 Hz) for H-6'. There is an AA'BB' system confirmed by the signals at  $\delta_{\text{H}}$  6.77 (d,  $J=8.4$  Hz) for H-3''', 5''' and  $\delta_{\text{H}}$  7.60 (2H, d,  $J=8.4$  Hz) for H-2''', 6'''. Acid hydrolysis of compound **6** yielded amentoflavone as the aglycone part which was confirmed by direct comparison to the isolated amentoflavone in this work on TLC. Paper chromatography of the sugar part resulting from acid hydrolysis showed that glucose is the glycone moiety. The glycosidic linkage was identified as  $\alpha$  as indicated by the signal at  $\delta_{\text{H}}$  4.47 ( $J=4.4$  Hz) for the anomeric proton. <sup>13</sup>C NMR signals at  $\delta_{\text{C}}$  99.9, 72.9, 70.2, 70.0, 67.9, 63.5 ppm confirmed the presence of one sugar moiety (*O*-glucose) and matched those for  $\alpha$  anomer<sup>[30]</sup>. The ESI<sup>+</sup>-MS of compound **6** showed an  $m/z$  at 701 for [M+H]<sup>+</sup>, which is a solid evidence for amentoflavone glucoside. The glucosidation of the 5, 5''- OH was ruled due to the presence of signals in the <sup>1</sup>H NMR spectrum for free hydroxyl groups at these sites as mentioned above and thus suggested the conjugation of either 7-OH, 7''-OH, 4'-OH or 4''-OH. <sup>1</sup>H NMR spectrum showed also signals at  $\delta_{\text{H}}$  10.30 (brs) and 10.85 (brs) which are assigned for 7- OH and 7''- OH, respectively<sup>[31]</sup>. However the signals for 4', 4''-OH were absent indicating possible conjugation at either position. Markham *et al* in 1978 indicated that glucosylation causes upfield shift of the *ipso* carbon and downfield shift of the *ortho* and *para* carbons, with greater impact on the *para* position, however only one example of glycosylation at C-4' was studied and the results were considered preliminary<sup>[26]</sup>. Literature of apigenin 4'-*O*- $\alpha$ -




Figure 1: Structures of compounds 1-8.

D-glucopyranoside indicated that the glycosylation has no significant impact on the *ipso*, *ortho* and *para* carbons<sup>[24,33]</sup> and thus the glycosylation rules are questionable at this site. <sup>13</sup>C NMR data of compound **6** showed that C-4' was shifted downfield (*ipso* carbon,  $\delta_C$  161.0) accompanied by downfield shift of the *ortho* carbons C-3', 5' ( $\delta_C$  121.9, 121.8, respectively), and the *para* carbon C-1' ( $\delta_C$  128.5) compared to the respective sites of amentoflavone isolated in this work ( $\delta_C$  159.6, 121.7, 116.4, and 120.5, respectively) and in literature<sup>[26,29]</sup>. However, there were no significant differences in the chemical shift of C-4'', C-3'', C-5'', and C-1'' resonating at  $\delta_C$  161.7, 116.7, 116.17, and 121.9, respectively compared to the respective sites in compound **5** observed at  $\delta_C$  161.5, 116.2, 116.2, 121.4, respectively and also to the literature data<sup>[29]</sup>. Thus, the <sup>13</sup>C NMR augments the conclusion to name compound **6** as amentoflavone-4'-O- $\alpha$ -D-glucopyranoside indicating that glycosylation of biflavonoids could behave differently from simple flavonoids (Figure 1). To our knowledge, this is the first report of monoglucosidated amentoflavone from plants elucidated by NMR and MS studies. Enzymatic synthesis of amentoflavone monoglucoside by Chaudhary

*et al* was accomplished by using recombinant oleandomycin glycosyltransferase<sup>[33]</sup>. However, the authors did not determine the site of glycosylation and only used LC-MS/MS for detection of the glycoside.

#### Compound 7

Compound **7** gave positive Molisch's test. UV spectrum showed  $\lambda_{max}$  at 283 and 340 (sh) nm, which suggested a flavonoid glycoside structure. <sup>1</sup>H-NMR spectrum showed the characteristic signals for flavanone structure represented by the signals at  $\delta_H$  2.55 (1H, dd,  $J=16.3, 3.2$  Hz) and 3.78 (1H, dd,  $J=16.3, 12.6$  Hz) for C-3 protons and  $\delta_H$  5.52 (1H, dd,  $J=12.6, 3.2$ ) for H-2. Signals for *meta*-coupled protons at  $\delta_H$  6.12 and 6.14 ( $J=3.6$  Hz) are consistent with signals due to H-6 and H-8. An ABX system was displayed by signals resonating at  $\delta_H$  6.96 (d,  $J=1.6$  Hz) for H-2', 6.94 (dd,  $J=1.6, 8.4$  Hz) for H-6' and 6.91 (d,  $J=8.4$  Hz) for H-5'. A signal at  $\delta_H$  3.9 integrated for 3 protons is consistent with a methoxy group. Signals for two anomeric protons resonating at  $\delta_H$  4.69 (d,  $J=2.8$  Hz) and 4.97 (d,  $J=7.4$  Hz) suggested the presence of  $\alpha$ -L-rhamnose and  $\beta$ -D-glucose. The presence of two hydroxyls

at C-3' and C-5 was evidenced by signals at  $\delta_H$  9.12 and 12.01, respectively, which indicated that compound **7** is an *O*-glycoside of a disaccharide at either C-7 or C-4'. The  $^{13}C$  NMR data established the site and the linkage of the disaccharide conjugation. The upfield shift of C-7 by 2.2 ppm and downfield shift of C-6, C-8 and C-10 by 1.7, 1.5 and 1.5 ppm, respectively in comparison with the aglycon data in literature confirmed the site of conjugation at C7<sup>[32, 34]</sup>. The inter-linkage of rhamnose and glucose is determined to be (1→2) and not (1→6), which was supported by the downfield shift of C-2 signal of glucose at  $\delta_C$  76.7 while C-6 was represented by the signal at  $\delta_C$  59.8 which is consistent with hesperidoside disaccharide<sup>[30]</sup>. ESI<sup>+</sup>-MS of compound **7** showed an [M+H]<sup>+</sup> peak at *m/z* 611 which is in concordance with the determined structure. The NMR and MS data of compound **7** confirmed the structure as neohesperidine which matches those in literature<sup>[23, 34-36]</sup>.

#### Compound 8

Compound **8** gave positive Molisch's test. UV spectrum showed  $\lambda_{max}$  at 260 and 357 nm which suggested a flavonoid glycoside structure. UV spectra of compound **8** with different shift reagents suggested a flavonol structure with free hydroxyls at positions 5, 7, 3' and 4'.  $^1H$ -NMR spectrum confirmed the flavonol structure by the absence of the H-3 singlet. Signals for *meta*-coupled protons at  $\delta_H$  6.16 and 6.33 (*J*= 2.0 Hz) are consistent with H-6 and H-8 assignment. Signals at  $\delta_H$  7.54 (m) and 6.83 (d, *J*= 8.4 Hz) were assigned for H-2', H-6' and H-5', respectively. Signals at  $\delta_H$  4.39 (d, *J*= 2.0 Hz) and 5.32 (d, *J*= 7.0 Hz) matches those in literature for  $\alpha$ -L-rhamnose and  $\beta$ -D-glucose, respectively<sup>[35]</sup>. Direct comparison of compound **8** with rutin authentic sample by TLC and superimposable IR alongside the literature data<sup>[37, 38]</sup> confirmed the structure as rutin.

#### BIOLOGICAL ACTIVITY

The antioxidant activity of the methanol extract and its fractions (pet. ether, methylene chloride, ethyl acetate and n-butanol) was investigated using DPPH method. Of the tested extracts, methylene chloride and ethyl acetate fractions gave the best results (Table 1). The activity of methylene chloride fraction can be attributed to its content of amentoflavone and naringenin. Amentoflavone is a well known antioxidant and free radical scavenger<sup>[39]</sup>. Naringenin exhibits substantial free radical scavenging and

DNA protection properties as shown by Cavia –Saiz *et al*<sup>[40]</sup>. The activity of the ethyl acetate fraction was investigated further by testing the activity of the pure compounds isolated from this fraction (amentoflavone glucoside **6** and neohesperidin **7**) alongside amentoflavone **5**. The SC<sub>50</sub> of amentoflavone glucoside and neohesperidin were found 14.8 and 13.2  $\mu$ g/ml, respectively which are lower than that of quercetin (18  $\mu$ g/ml). Amentoflavone exhibited an SC<sub>50</sub> of 24.2  $\mu$ g/ml. The results indicated that neohesperidin and amentoflavone glucoside are potent free radical scavengers, three to four times more potent than quercetin at low dose levels (6.25  $\mu$ g/ml) (Table 2) and even more potent than amentoflavone itself.

Cytotoxic activity of the methanol extract and its fractions (pet. ether, methylene chloride, ethyl acetate and n-butanol) against MCF7 breast cancer cell and HepG2 liver cancer cell line was investigated using SRB assay. Ethyl acetate fraction showed the highest inhibitory activity (69.06%) against MCF7 cell line followed by n-butanol and methylene chloride fractions (66.58% and 65.93%, respectively) (Table 3). n-Butanol fraction was the most active extract against HepG2 cell line (68.4%) followed by pet. ether and methylene chloride (64.12% and 62.15%, respectively). The percentage inhibition of n-butanol fraction against HepG2 cell line can be justified by its rutin content. Cristina *et al* investigated the cytotoxicity and protective activity of rutin against the pro-carcinogen benzo(a)pyrene and their results suggested that rutin is a cytotoxic agent against HTC hepatic cells and has a protective effect against DNA damage<sup>[41]</sup>. Naringenin and amentoflavone possess anticancer action,<sup>[42, 43]</sup> which can account for the cytotoxic activity displayed by the methylene chloride fraction in our study. Amentoflavone alongside amentoflavone glucoside and neohesperidin were tested for their cytotoxic activity against MCF7 cell lines to explain the activity of the ethyl acetate fraction. Neohesperidin showed an IC<sub>50</sub> of 4.73  $\mu$ g/ml, which is comparable to that of doxorubicin (IC<sub>50</sub> of 4.13  $\mu$ g/ml) followed by amentoflavone glucoside (IC<sub>50</sub> of 6.12  $\mu$ g/ml), whilst amentoflavone exhibited an IC<sub>50</sub> of 18.70  $\mu$ g/ml (Table 4). Our results of antioxidant and cytotoxic activity investigation demonstrated that amentoflavone glucoside and neohesperidin are promising antioxidant and cytotoxic agents which awaits further studies towards understanding the molecular mechanism of their action.

**Table 1: Antioxidant activity of different extracts**

Sample	% Scavenging activity $\pm$ SE				
	Conc. (mg/ml)				
	10	7.5	5	2.5	1.25
Quercetin	100	100	100	97.5 $\pm$ 0.03	76.3 $\pm$ 0.04
Alcohol extract	64 $\pm$ 0.07	60.7 $\pm$ 0.03	58.9 $\pm$ 0.03	44.8 $\pm$ 0.03	38.4 $\pm$ 0.03
Pet. ether extract	69.2 $\pm$ 0.05	63.5 $\pm$ 0.04	59.1 $\pm$ 0.05	54 $\pm$ 0.02	39.4 $\pm$ 0.04
Methylene chloride extract	74.2 $\pm$ 0.05	68.5 $\pm$ 0.05	66.5 $\pm$ 0.04	52.5 $\pm$ 0.06	43.3 $\pm$ 0.05
Ethyl acetate extract	74.2 $\pm$ 0.04	73.3 $\pm$ 0.03	70.1 $\pm$ 0.04	52.5 $\pm$ 0.05	47.1 $\pm$ 0.03
n-Butanol	63.4 $\pm$ 0.04	56.3 $\pm$ 0.03	55.1 $\pm$ 0.05	48.8 $\pm$ 0.04	46.4 $\pm$ 0.05

SE= standard error, n=3

**Table 2: Antioxidant activity of compounds 5, 6 and 7.**

Compound	% Scavenging activity $\pm$ SE					
	Conc. ( $\mu$ g/ml)					
	100	50	25	12.5	6.25	SC <sub>50</sub>
Quercetin	77.0 $\pm$ 0.05	76.1 $\pm$ 0.03	65.7 $\pm$ 0.04	30.0 $\pm$ 0.02	10 $\pm$ 0.04	18 $\pm$ 0.05
Amentoflavone	60.9 $\pm$ 0.03	60.1 $\pm$ 0.05	59.3 $\pm$ 0.05	33 $\pm$ 0.02	19.4 $\pm$ 0.05	24.2 $\pm$ 0.06
Amentoflavone glucoside	66.2 $\pm$ 0.02	63.8 $\pm$ 0.03	62.5 $\pm$ 0.052	45.3 $\pm$ 0.06	43.2 $\pm$ 0.02	14.8 $\pm$ 0.05
Neohesperidin	64.5 $\pm$ 0.03	63.1 $\pm$ 0.05	60.9 $\pm$ 0.04	45.3 $\pm$ 0.02	33.4 $\pm$ 0.04	13.2 $\pm$ 0.05

SE= standard error, n=3

**Table 3: Cytotoxic activity of the tested extracts using single dose (100  $\mu$ g/ml)**

Extract	% Inhibition $\pm$ SE	
	MCF7	HepG2
	Alcohol	62.16 $\pm$ 0.4
Pet. ether	63.89 $\pm$ 0.22	64.12 $\pm$ 0.12
Methylene chloride	65.93 $\pm$ 0.18	62.15 $\pm$ 0.14
Ethyl acetate	69.06 $\pm$ 0.14	59.85 $\pm$ 0.18
n-Butanol	66.58 $\pm$ 0.14	68.4 $\pm$ 0.18

SE: standard error, n=3

**Table 4: Cytotoxic activity of compounds 5, 6 and 7 using MCF-7 cell line.**

Compound	Surviving fraction $\pm$ SE				
	Conc. ( $\mu$ g/ml)				
	5	12.5	25	50	IC <sub>50</sub>
Doxorubicin	0.40 $\pm$ 0.031	0.44 $\pm$ 0.029	0.37 $\pm$ 0.035	0.40 $\pm$ 0.031	4.13
Amentoflavone	0.75 $\pm$ 0.014	0.58 $\pm$ 0.036	0.42 $\pm$ 0.016	0.43 $\pm$ 0.037	18.70
Amentoflavone glucoside	0.52 $\pm$ 0.051	0.32 $\pm$ 0.029	0.22 $\pm$ 0.018	0.20 $\pm$ 0.016	6.12
Neohesperidin	0.49 $\pm$ 0.013	0.51 $\pm$ 0.021	0.44 $\pm$ 0.025	0.41 $\pm$ 0.029	4.73

SE: Standard error, n=3.

### CONCLUSION

Chemical investigation of *C. revoluta* leaflets indicated that flavonoids and biflavonoids are the main principals. A new amentoflavone glucoside was reported for the first time in this paper based on NMR and MS analyses.  $\beta$  amyryn (**1**), neohesperidin (**7**), and rutin (**8**) were reported for the first time from *Cycas* genus.  $\beta$  sitosterol (**2**) was isolated for the first time from *C. revoluta* Thunb.

Biological screening of the isolated compounds showed an interesting activity profile for amentoflavone-4'-*O*- $\alpha$ -D-glucopyranoside and neohesperidin as antioxidant and cytotoxic agents which requires further studies for better understanding and possible harnessing in medicine.

### REFERENCES

- Rani MS, Rao CV, Gunasekar D, Blond A, Bodo B.A. Biflavonoids from *Cycas beddomei*. *Phytochemistry* 1998;47:319-21.
- Duke JA. CRC handbook of medicinal herbs. 2<sup>nd</sup> edition Boca Raton, Florida: CRC Press;1985. p. 643-44.
- Walters T, Osborne R. Cycad classification concepts and Recommendation CABI publishing, Walling ford UK; 2004. p. 1-11.
- Jones D. Cycads of the World. Washington D. C: Smithsonian Institution Press; 1993.
- Moawad A, Hetta M, Zjawiony JK, Ferreira D, Hifnawy M. Two new dihydroamentoflavone glycosides from *Cycas revoluta*. *Nat Prod Res* 2014;28(1): 41-7.
- Moawad A, Hetta M, Zjawiony JK, Tacob MR, Hifnawy M, Ferreira D. Phytochemical investigation of *C. circinalis* and *C. revoluta*. *Planta Med* 2010; 76(8):796-2.
- Mourya MK, Prakash A, Swawi A, Singh GK, Mathura. Leaves of *Cycas revoluta*, potent antimicrobial and antioxidant agent. *World J Sci Technol* 2011;1(10):11-20.
- Duke JA, Ayensu ES. Medicinal plants of China. 1<sup>st</sup> edition. Algonac, Michigan: Reference Publication;1985.
- Varshney AK, Mah T, Khan NU, Rahman W, Hwa CW, Okigawa M *et al*. Biflavonoids from *Cycas revoluta*, *C. circinalis* and *C. rumphii*. *Indian J Chem* 1973;11:1209-14.
- El said ME, Amer MM. Oils, fats, waxes and surfactants. Anglo-Egyptian Book shop. Cairo;1965.
- Harborne JB, Mabry TJ, Mabry H. The flavonoids. Academic Press, New York, San Fransisco. 1975. p. 387.
- Kabbash A; studies on anti-oxidative natural products and bioactive ingredients in Aloe Vera, Ph. D Thesis, Faculty of Pharmacy, Tanta University 2002;27-30.
- Gulluce M, Sahin F, Sokmen M, Ozer H, Daferera A. Antimicrobial and antioxidant properties of the essential oils methanol extract from *Mentha longifolia* L. sp. *Longifolia*. *Food Chem* 2007;103(4):1449-56.
- Stanojevic L, Stankovic M, Nikolic V, Nikolic L, Ristic D *et al*. Antioxidant activity and total phenolic Flavonoid Contents of *Hieracium pilosella* L. extracts. *Sensors* 2009;9(7):5702-14.
- Skehan P, Strohreng R, Scudiero D, Monks A, Mahon J, Vistica D *et al*. New colorimetric cytotoxicity assay for anticancer-drug screening. *J Natl Cancer Inst* 1990;82(13):1107-12.
- Ayyad SE, Abdel-Lateff A, Alarif WM, Patacchioli FR, Badria FA, Ezmirly ST. In vitro and in vivo study of Cucurbitacins-type triterpene glucoside from *Citrullus colocynthis* growing in Saudi Arabia against hepatocellular carcinoma. *Environ Toxicol Pharmacol*. 2012; 33(2): 245-51.
- Vázquez LH, Palazon J, Ocaña AN. The Pentacyclic Triterpenes  $\alpha$ ,  $\beta$ -amyryns: A review of sources and biological activities. *Phytochemicals- A Global Perspective of Their Role in Nutrition and Health*. 2012. p. 487-502.

18. Trivedi PC, Chaudhery N. Isolation and characterization of bioactive compound  $\beta$ -sitosterol from *Withania somnifera* L. JPR 2011;4(11):4252-53.
19. Khanam S and Sultana R. Isolation of  $\beta$ -sitosterol and stigmasterol as active immunomodulatory constituent from the fruits of *Solanum xanthocarpum* (Solanaceae). IJPSR 2012;3(4):1057-60.
20. Sen A, Dhavan P, Kshitiz, Singh KK, Tejavathi G. Analysis of IR, NMR and anti-microbial activity of  $\beta$ - Sitosterol isolated from *Momordica charantia*. SSJBt 2013;1(1):9-13.
21. Ishii K, Furuta T, Kasuya Y. Determination of naringin and naringenin in human plasma by high-performance liquid chromatography. J Chromatogr B Biomed 1996;683(2):225-29.
22. Shafaghat A, Salimi F. Extraction and Determining of Chemical Structure of Flavonoids in *Tanacetum parthenium* (L.) Schultz. Bip. from Iran. JSIAU 2008;18( 68):39-42.
23. Maltese F, Erkelens C, Kooy F, Hae Choi Y, Verpoorte R. Identification of natural epimeric flavanone glycoside by NMR spectroscopy. Food chemistry J 2009;116(2):575-79.
24. Zhang YM, Zhang YM, Tan NH, Huang HO, Jia RR, Zeng GZ, Ji CJ. Three bioactive biflavone isolated from *Taxodium mucronatum*. Acta Bot Yunnan 2005;27(1):107-10.
25. Markham KR. The structures of amentoflavone glycosides isolated from *Psilotum nudum*. Phytochemistry 1984;23(9):2053-56.
26. Markham KR, Geger H, Mabry J. Carbon-13 NMR studies of Flavonoids-III naturally occurring flavonoid glycosides and their acylated derivatives. Tetrahedron 1978;34(9):1389-98.
27. Andersen M, Markham KR. Flavonoids Chemistry, Biochemistry and Applications. Taylor and Francis, Boca Raton, USA. 2006. p. 57-67.
28. Harborne JB, Mabry TJ, Mabry H. The flavonoids. 1<sup>st</sup> ed. Champan and Hall Ltd. London; 1975; 2: 46- 61.
29. Markham KR, Shepard C, Geiger H. <sup>13</sup>C NMR studies of some naturally occurring amentoflavone and hinokiflavone biflavonoids. Phytochemistry 1987;26(12):3335-37.
30. Agrawal PK. Carbone -13NMR of flavonoids. ELSIEVIER. Amsterdam. Oxford. New York. Tokyo. 1989. p. 274-89.
31. Jinga Y, Zhanga G, Mab E, Zhanga H, Guanc J, He J. Amentoflavone and the extracts from *Selaginell tamariscina* and their anticancer activity. Asian J Tradit Med 2010;5(6):226-29.
32. Gaimei S, Zhiqin G, Haining L, Dongm S. New flavonoid glycoside from *Elsholtzia rugulosa* Hemsl. Molecules 2009;14(10):4190-96.
33. Chaudhary AK, Hwang IY, Jo YJ, Choi SH, Lee EY. Enzymatic synthesis of amentoflavone glycoside using recombinant oleandomycin glycosyltransferase. J Ind Eng Chem 2015; 25: 304-307.
34. Erenler R, Sen O, Akisit H, Demirtas I, Yaglioglu AS, Elmastas M *et al.* Isolation and identification of chemical constituents from *Origanum majorana* and investigation of antiproliferative and antioxidant activities. Jsfa 2015; doi:10.1002/ Jsfa.7155.
35. Ternai B, Markhan KR. <sup>13</sup>C NMR of flavonoids. Tetrahedron 1976;32:2607- 12.
36. Kometani T, Nishimura T, Nakae T, Takii H, Okada S. Synthesis of neohesperidin glycosides and naringin glycosides by cyclodextrin glucanotransferase from an alkalophilic Bacillus Species. Biosci Biotechnol Biochem 2006;60 (4).645-49.
37. Sintayehu B, Asres K, RaghavendraY. Radical scavenging activities of the leaf extracts and a flavonoid glycoside isolated from *Cineraria abyssinica* Sch. Bip. Exa. Rich. JAPS 2012;2(4):44-9.
38. Klimek B, Tokar M. Biologically active compounds from the flowers of *Forsythia susensa* vahl. Acta Poloniae pharmaceutica-Drug Research 1998; 55(6):499-04.
39. Pawlak K, Bylka W, Jazurek B, Matlawska I, Sikorska M, Manikowski E *et al.*G. Antioxidant activity of flavonoids of different polarity assayed by modified ABTS cation radical decolorization and EPR Technique. Acta Biologica Cracoviensia series Botanica 2010;52(1):97-04.
40. Cavia-Saiz M, Busto MD, Pilar-Izquierdo MC, Ortega N, Perez-Mateos M, Muñiz P. Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin, a comparative study. J Sci Food Agric 2010;90(7):1238-44.
41. Marcarini JC, Tsuboy MS, Luiz RC, Ribeiro LR, Hoffmann CC, Mantovani MS. Investigation of cytotoxic, apoptosis-inducing, genotoxic and protective effects of the flavonoid rutin in HTC hepatic cells. Exp Toxicol Pathol 2011;63(5):459-65.
42. Setyawan A.D. Natural products from genus Selaginella (Selaginellaceae). Nus Biosci 2011; 3(1):44-58.
43. Zhang FY, Du GJ, Zhang L, Zhang CL, Lu WL, Liang W. Naringenin enhances the anti-tumor effect of doxorubicin through selectively inhibiting the activity of multidrug resistance-associated proteins but not P-glycoprotein. Pharm Res 2009;26(4):914-25.