































- [266]. Rossi, E.J., Sim, L., Kuntz, D.A., Hahn Johnshen, B.D., Ghavami, A., Szczepiana, M.G., Kumar, N.S., Sterchi, E.E., Nicholas, B.L., Pinto, B.M., and Rose, D.R. Inhibition of recombinant human maltase glucomylase by salacinol & derivatives. *FEBS J.* 2006; 273: 2673-2683.
- [267]. Huang, T.H.W., Yang, Q., Harada, M., Uberai, J., Radford, J., Li, G.Q., Yamahara, J., Roufogalis, B.D., Li, Y. *Salacia oblonga* root improves cardiac lipid metabolism in Zucker diabetic fatty rats: modulation of cardiac PPAR- $\alpha$ -mediated transcription of fatty acid metabolic genes. *Toxicol. App. Pharmacol.* 2006; 210: 78-85.
- [268]. Grover, J.K., Vats, V., Rathi, S.S. and Dawar, R. Traditional Indian anti-diabetic plants attenuate progression of renal damage in streptozotocin induced diabetic mice. *J. Ethnopharmacol.* 2001; 76: 233-238.
- [269]. Welles, B. Glucocorticoids in Type 2 Diabetes Mellitus and the metabolic syndrome. *Drug Dev. Res.* 2006; 67: 570-573.
- [270]. Huang, T.H.W., He, L., Qin, Q., Yang, Q., Peng, G., Harada, M., Qi, Y., Yamahara, J., Roufogalis, B.D. and Li, Y. *Salacia oblonga* root decreases cardiac hypertrophy in Zucker diabetic fatty rats: inhibition of cardiac expression of angiotensin II type 1 receptor. *Diabetes Obes. Metab.* 2008; 10 (7): 574-585.