









- [24] Marinho VC, Higgins JP, Sheiham A, Logan S. Fluoride toothpastes for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* 2003;(1):CD002278 Epub 2003/01/22. doi: 10.1002/14651858.CD002278 . [PubMed] [Ref list]
- [25] Santos AP, Oliveira BH, Nadanovsky P. Effects of low and standard fluoride toothpastes on caries and fluorosis: systematic review and meta-analysis. *Caries Res.* 2013;47(5):382–90. Epub 2013/04/11.
- [26] fluoridealert.org > issues > dental-products
- [27] Mishra V, Shettar L, Bajaj M, Math AS. Comparison of a commercially available herbal and 0.2% chlorhexidine mouthrinse for prevention of oral malodor: A clinical trial. *Journal of International Society of Preventive & Community Dentistry.* 2016;6(Suppl 1):S6-S11. doi:10.4103/2231-0762.181160.
- [28] Yousefimanesh H, Amin M, Robati M, Goodarzi H, Otoufi M. Comparison of the antibacterial properties of three mouthwashes containing chlorhexidine against oral microbial plaques: An in vitro study. *Jundishapur J Microbiol.* 2015;8:e17341. [PMC free article] [PubMed]
- [29] Rosenberg M, Gelernter I, Barki M, Bar-Ness R. Day-long reduction of oral malodor by a two-phase oil:water mouthrinse as compared to chlorhexidine and placebo rinses. *J Periodontol.* 1992;63:39–43. [PubMed]
- [30] Supranoto SC, Slot DE, Addy M, Weijden GA. The effect of CHX dentifrice or gel versus CHX mouthwash on plaque, gingivitis, bleeding, tooth discoloration: A systematic review. *Int J Dent Hyg.* 2015;13:83–92.