



















29. Wang, X., Jiang, C., Ko, T. et al. Correlation between optic disc perfusion and glaucomatous severity in patients with open-angle glaucoma: An optical coherence tomography angiography study. *Graefes Arch Clin Exp Ophthalmol*, 2015; **253** (9): 1557-1564.
30. Wang, Y., Bower, D., Izatt, J., Tan, O. and Huang, D. Retinal blood flow measurement by circumpapillary Fourier domain Doppler optical coherence tomography. *J Biomed Opt*, 2008; **13** (6): 064003. doi: 10.1117/1.2998480.
31. Wang, Y., Fawzi, A. A. and Varma, R. et al. Pilot study of optical coherence tomography measurement of retinal blood flow in retinal and optic nerve diseases. *Invest Ophthalmol Vis Sci*, 2015; **52**: 840-845.
32. Weinreb, R.N. and Harris, A. *Ocular Blood Flow in Glaucoma: Consensus Series 6*. The Netherlands: Kugler Publications, 2007.
33. Yaoeda, K. and Shirakashi, M. et al. Relationship between optic nerve head circulation and visual field loss in glaucoma. *Acta Ophthalmol Scand*, 2013; **81**: 253.
34. Yarmohammadi, A., Zangwill, L. M. and Diniz-Filho, A. et al. Optical coherence tomography angiography vessel density in healthy, glaucoma suspect, and glaucoma eyes. *Invest Ophthalmol Vis Sci*, 2016; **57** (9): 451-459.
35. Yu, J., Gu, R., Zong, Y. et al. Relationship between retinal perfusion and retinal thickness in healthy subjects: an optical coherence tomography angiography study. *Invest Ophthalmol Vis Sci*, 2016; **57**: 204-210.