

REFERENCES

- Riaz MN, *Extruders in Food Applications* (2nd edn). CRC Press ,Florida, United States, 2000, pp.1-240
- Guy R, *Extrusion Cooking: Technology and Applications*, CRC Press ,Woodhead Publishing Limited (8th edn).Cambridge,England,2001,pp.1-212 .
- Fellows PJ, *Food Processing Technology: Principles and Practice* (4th edn), Woodhead Publishing Series in Food Science, Technology and Nutrition,2016, pp.753-774 .
- Aylin Altan, Kathryn LM and Medeni Maskan, Extrusion cooking of barley flour and process parameter optimization by using response surface methodology, *Journal of the Science of Food and Agriculture* 2008; **88**:1648–1659.
- Serge EO, Gu BJ, Kim YS and Ryu GH, Effects of feed moisture and barrel temperature on physical and pasting properties of cassava starch extrudate, *Korean J Food Preserv* 2011; **18** : 271–278.
- Iwe MO, Effects of extrusion cooking functional properties of mixtures of full-fat soy and sweet potato, *Plants Foods and Human Nutrition* 1998; **53** : 37-46.
- Bravo L, Siddhuraju P and Saura-Calixto F, Composition of underexploited Indian pulses. Comparison with common legumes. *Food Chemistry* 1999; **64**: 185–192.
- Ray PK, Toxic factor(s) in raw horse gram (*Dolichos biflorus*),*Journal of Food Science* 1969; **6**:207–211.
- Sreerama YN, Sasikala VB and Pratapa VM, Nutritional implications and flour functionality of popped/expanded horse gram, *Food Chemistry* 2008; **108**: 891–899.
- Geervani P and Eggum BO, Nutrient composition and protein quality of minor millets. *Plants Foods and Human Nutrition* 1989; **39**: 201–208(1989).
- Saleh AA and Tarek AE,Nutritional composition of chickpea (*Cicer arietinum* L.) as affected by microwave cooking and other traditional cooking methods. *J Food Comp Anal* 2006; **19**: 806–812.
- Geetha R, Mishra HN and Srivastav PP, Twin screw extrusion of kodo millet-chickpea blend: process parameter optimization, physico-chemical and functional properties, *Journal of Food Science and Technology* 2012; **51**: 44-53.
- Lai LS and Kokini JL,Physicochemical Changes and Rheological Properties of Starch during Extrusion.*Biotechnology Progress* 1991; **7**:251-266.
- Sibel Yagcı and Fahrettin Gogus, Response surface methodology for evaluation of physical and functional properties of extruded snack foods developed from food-by-products. *Journal of Food Engineering* 2008; **86** : 122–132.
- Emmanuel Kwasi Asare, Peluola-Adeyemi, Micheal Ayodeleldowo, Lateef Oladimeji Sanni and Goke Jacob Bodunde, Application of response surface methodology for studying the product characteristics of extruded rice/cowpea/groundnut blends. *International Journal of Food Sciences and Nutrition* 2004; **55** : 431 -439.
- Oluwaranti Abiodun, Peluola Adeyemi, MichealAyodeleldowo, Lateef Oladimeji Sanni and Goke Jacob Bodunde, Effect of extrusion parameter on the nutrient composition and quality of snacks developed from cocoyam (*Xanthosoma sagittifolium*) flour.*African Journal Of Food Science* 2014; **8**: 510-518 .
- Chakraborty SK, Singh DS, Kumbhar BK and Singh D. Process parameter optimization for textural properties of ready-to-eat extruded snack food from millet and legume pieces blends. *Journal Texture Studies* 2009; **40** : 710-726.
- Renu Sharma,Raj kumarTanuja srivastava, Saxena DC. (2015). Textural and Microstructural Properties of Extruded Snack Prepared from Rice Flour, Corn Flour and Deoiled Rice Bran by Twin Screw Extrusion, International Conference on Advancements in Engineering and Technology 2015, pp.0975 – 8887.
- AOAC, *Official Methods of Analysis of the AOAC* (16th edn). Association of Official Analytical Chemists, Washington DC,1990.
- AOAC. *Official Methods of Analysis of the AOAC* (18th edn). Association of Official Analytical Chemists, Gaithersburg, MD,2005.
- AOAC. *Official Methods of Analysis of the AOAC* (18th edn). Association of Official Analytical Chemists, Gaithersburg, MD,1995.
- Lowry OH, Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 1951; **193** : 265–275.
- Ying Gui, Sun Kuk Gil and Gi Hyung Ryu, Effects of Extrusion Conditions on the Physicochemical Properties of Extruded Red Ginseng. *Preventive Nutrition and Food Science* 2012; **17** : 203–209.
- Singleton VL and Rossi JA , Colorimetry of total phenolics with phosphor-molybdicphosphotungstic acid reagents. *American Journal for Enology and Viticulture* 1965; **16**: 144–58.
- Balunkeshwar nayak, Berrios JJ, Powers JR and Tang J, Effect of Extrusion on the Antioxidant Capacity and Color Attributes of Expanded Extrudates Prepared from Purple Potato and Yellow Pea Flour Mixes. *Journal of Food Science* 2011; **76** : 874-83.
- Flores-Morales A, Determination of the structural changes by FT-IR, Raman and CP/MAS 13C NMR spectroscopy on retrograded starch of maize tortillas. *Carbohydrate Polymers* 2012; **87** : 61-68.
- Carine Semasaka, Xiangzhen Kong and Yufei Hua, Optimization of Extrusion on Blend Flour Composed of Corn, Millet and Soybean. *Pakistan Journal Of Nutrition* 2010; **9** : 291-297.
- Chakraborty SK, Singh DS, Kumbhar BK and Singh D, Process parameter optimization for textural properties of ready-to-eat extruded snack food from millet and legume pieces blends. *Journal Texture Studies* 2009; **40** : 710-726.
- Shalini GR, Neetu Jakhar, Jyoti Nishad, Neha Saini, Sangita Sen, Rakesh Bhardhwaj, SarikaJaiswal and Poonam Suneja, Extrusion Conditions and Antioxidant Properties of Sorghum,Barley and Horse Gram Based Snack. *International Journal of Plant Research* 2015; **28** : 171-182.
- Singh D, Girraj Singh Chauhan, Suresh I and Tyagi SM, Nutritional quality of extruded snakes developed from composite of rice broken and wheat bran. *International Journal of Food Properties* 2000; **3** : 421-431.
- Deshpande HW and Poshadri A, Physical and sensory characteristics of extruded snacks prepared from Foxtail millet based composite flours. *International Food Research Journal* 2011; **18**: 751-756.
- Shivani Pathania, Baljit Singh, Savita Sharma, Vandana Sharma and Smita Singla, Optimization of extrusion processing conditions for preparation of an instant grain base for use in weaning foods. *International Journal of Engineering Research and Applications* 2013; **3**: 1040-1049.
- Oke MO, Awonorin SO, Sanni LO, Asiedu R and Aiyedun PO, Effect of varieties on physicochemical and pasting characteristics of water yam flours and starches. *Journal Of Food Processing And Preservation* 2012; **8**: 1745-4549.
- Sawant AA, Application Of Horse Gram Flour In Extruded Products. *International Journal of Development Research* 2015; **5**: 5668-5673.
- Basediya AL, Sheela Pandey, Shrivastava SP, Khurshed Alam Khan and Anura Nema, Effect of process and machine parameters on physical properties of extrudate during extrusion cooking of sorghum, horse gram and defatted soy flour blends. *Journal of Food Science and Technology* 2013; **50**: 44–52.
- Dogan H and Karwe MV, Physico-chemical Properties of quinoa extrudates, *Food Science and Technology International* 2003; **9**: 101-114.
- Martínez MM, Rosellb CM and Manuel Gómeza, Modification of wheat flour functionality and digestibility through different extrusion conditions. *Journal of Food Engineering* 2014; **143**: 74-79.
- Rodríguez-Miranda J, Ruiz-López II, Herman Lara E, Martínez Sánchez CE, Delgado LE and Vivar Vera MA, Development of extruded snacks using taro (*Colocasiaesculenta*) and nixtamalized maize (*Zea mays*) flour blends. *Food Science and Technology* 2011; **44**: 673-680.
- Bourne, MC. Food texture and viscosity: Concept and measurement. Academic Press, San Diego, CA. edn2,1982, p. 256.
- Owusu Anshah J, Van de Voort,FR,Stanley,DW. Physicochemical changes in cornstarch as a function of Extrusion variables,Cereal chemistry 1984; **60**(4): 319-324.
- Yaqoub AG, Mohammed MA and Abu Baker AA, Effect of soaking, sprouting and cooking on chemical composition, bioavailability of minerals, vitamins and protein digestibility of Roselle (*Hibiscus sabdariffa* L.) seed. *Pakistan Journal of Nutrition* 2008; **7**: 50-56.
- Anuonye JC, John Onuh, Evans Egwim and Adeyemo SO, Nutrient and antinutrient composition of extruded acha/soybean blend. *Journal Of Food Processing And Preservation* 2010; **34**: 680-691 (2010).
- Guzmán-Ortiz FA, Hernández-Sánchez H, Yee-Madeira H, Martín-Martínez ES, María del Robles-Ramírez C, Rojas-López M, Jose Berrios DJ and Mora-Escobedo R, Physico-chemical, nutritional and infrared spectroscopy evaluation of an optimized soybean/corn flour extrudate. *Journal of Food Science and Technology* 2015; **52** : 4066–4077.