























- 37 Nakayama T, Watanabe M, Teramoto T, Kitajima M. CA19-9 as a predictor of recurrence in patients with colorectal cancer. *J Surg Oncol* 1997;66: 238-43
- 38 Barbara C. Wolf, John C. D'Emilia, Ronald R. Salem, Deborah DeCoste, Henry F. Sears, Leonard S. Gottlieb, Glenn D. Steele, Jr. Detection of the Tumor-Associated Glycoprotein Antigen (TAG-72) in Premalignant Lesions of the Colon. *Natl Cancer Inst* 2015; 81 (24): 1913-1917.
- 39 Umar SB, DiBaise JK. Protein-losing enteropathy: case illustrations and clinical review. *Am J Gastroenterol.* 2010;105:43-9.40-Lindmark G, Kressner U, Bergstrom R, Glimelius B. Limited clinical significance of the serum tumour marker CA 72-4 in colorectal cancer. *Anticancer Res* 1996;16:895-8.
- 40 American Association for Cancer Research(2010).
- 41 Xavier Filella, Jose Fuster, Rafael Molina, Juan Jose Grau, Luis Grande & Antonio M. Ballesta. Tag-72, CA 19.9 and CEA as Tumor Markers in Gastric Cancer; *Acta Oncologica* 1994;Vol. 33, No. 7, pp. 747-751.
- 42 Huang ZB, Zhou X, Xu J, Du YP, Zhu W, Wang J. Prognostic value of preoperative serum tumor markers in gastric cancer. *World J Clin Oncol* 2014;5:170-6.
- 43 Fiorella Guadagni, Mario Roselli, Maurizio Cosimelli, Ernest Mannella, Manfred Tedesco, Francesco Cavaliere, Antonio Grassi, Maria Rosaria Abbolito, Ph.D, John W. Greiner, PhD, and Jeffrey Schlo. TAG-72 (CA 72-4 Assay) as a Complementary Serum Tumor Antigen to Carcinoembryonic Antigen in Monitoring Patients with Colorectal Cancer 1991; Grant A.I.R.C.
- 44 Mattar R, Alves de Andrade CR, DiFavero GM, Gama-Rodrigues JJ, Laudanna AA. Preoperative serum levels of CA 72-4, CEA, CA 19-9, and alpha-fetoprotein in patients with gastric cancer. *Rev Hosp Clin Fac Med Sao Paulo* 2002; 57: 89-92.
- 45 Zheng CX, Zhan WH, Zhao JZ, Zheng D, Wang DP, He YL, Zheng ZQ. The prognostic value of preoperative serum levels of CEA, CA19-9 and CA72-4 in patients with colorectal cancer. *World J Gastroenterol* 2001;7(3):431-434.
- 46 Ychou M, Tuszinski T, Pignon JP, et al. Adenocarcinomes gastriques: comparaison du CA 19-9 et de l'antigène carcinoembryonnaire pour le diagnostic des récurrences après traitement chirurgical. *Gastroenterol Clin Biol*; 1992; 16: 848-52.
- 47 Guadagni F, MD, Mario Roselli, Teresa Aamto, Maurizio Cosimelli, Ernest Mannella, MD, Manfred Tedesco, Antonio Grassi, Vincenzo Casale, MD, Francesco Cavaliere, John W. Greiner, and Jeffrey Schlom(1991), Clinical Evaluation of Serum Tumor-Associated Glycoprotein-72 as a Novel Tumor Marker for Colorectal Cancer Patients; *Journal of Surgical Oncology Supplement* 2: 16-20.
- 48 Ebrahim Eftekhari, Hajar Jaberie, Fakhraddin Naghbalhossaini(2016).
- 49 Fiorella Guadagni, Mario Roselli, Maurizio Cosimelli, Antonella Spila, Francesco Cavaliere, Raffaella Arcuri, Maria Rosaria Abbolito, John W. Greiner, Jeffrey Schlom,. Biologic Evaluation of Tumor-Associated Glycoprotein-72 and Carcinoembryonic Antigen Expression in Colorectal Cancer, Part I 1994; volume. 37 no.2 .
- 50 Wanebo HJ, Rao B, Pinsky CM, Hoffman RG, Stearns M, Schwartz MK, Oettger HF. Preoperative carcinoembryonic antigen levels as a prognostic indicator of colorectal cancer. *N Engl J* 1978; Med 299:448-45.
- 51 Magdalena Świdarska, Barbara Choromańska, Ewelina Dąbrowska, Emilia Konarzewska-Duchnowska, Katarzyna Choromańska, Grzegorz Szczurko, Piotr Myśliwiec, Jacek Dadan, Jerzy Robert Ładny, Krzysztof Zwierz. The diagnostics of colorectal cancer ;*Contemp Oncol (Pozn)* 2014;18 (1): 1-6