


like candida will cause mucosal ulceration or even chronic inflammation. M/F RATIO

Off the positive patients



Though this study is conducted in the diabetic patients to know about the susceptibility to candidial infection, the other predisposing factors are all to be considered for their susceptibility. The amount of dietary carbohydrate and the other endocrine functions shall contribute more. Candidial infections are more pronounced in individuals with diminished cell mediated immunity, which is common feature in diabetes. Some time it will be seen as a consequence of alteration in the oral flora during long term antibiotic therapy or with broad spectrum antibiotic. Candidiasis is seen in during pregnancy but many women suffer with recurring episodes of candidiasis.



Image 1



Image 2

CONCLUSION

This study is conducted with the aim of assessing the health status of the diabetic patients. Presence of candida is taken as an indicator of the compromised state in diabetic patients. These patients are susceptible to any infections when their immune status is compromised. Candida is known to cause many types of oral lesion in patients attending dental clinics. It predominates in diabetic patients

REFERENCE

- 1- Manfredi M, Al-Karaawi Z, McCullough MJ, Hurel S, Porter SR. The isolation, identification and molecular analysis of Candida spp. isolated from the oral cavities of patients with diabetes mellitus. *Oral Microbiol Immunol* 2002; 17: 181-185
- 2- Fahad Ali Alshehri, Fawad Javed; Oral candida carriage and species prevalence among tobacco smokers and non smokers with and without type 2 diabetics; *OHDm-VOL 14*; February 2015
- 3- BV Kumar, NS Padshetty, KY Bai, MS Rao; prevalence of candida in the oral cavity of diabetics subjects; *JAPI VOL-53*; July 2005
- 4- Khaled H Abu-Elteen, Mawieh A Hamad, Suleiman A. Salah; prevalence of oral candida infection in diabetic patients; *Bahrain Medical Bulletin*; Vol-28 march 2006